arXiv:0905.2415v1 [gr-qc] 15 May 2009

On multitemporal generalization of Newton's gravitational law

V.D. Ivashchuk[1]

Center for Gravitation and Fundamental Metrology, VNIIMS, 46 Ozyornaya ul., Moscow 119361, Russia
Institute of Gravitation and Cosmology, Peoples' Friendship University of Russia, 6 Miklukho-Maklaya ul., Moscow 117198, Russia

Abstract

A n-time generalization of Newton's law (of universal gravitation) formula in $N = n + d + 1$-dimensional space-time is conjectured. This formula implies a relation for effective N-dimensional gravitational constant $G_{eff} = Gcos^2\theta$, where θ is the angle between the direction of motion of two particles in n-dimensional time manifold $\mathbb{R}^n$.

1 Introduction

In [1] a n-time generalization of the Newton's gravitational law formula in the $N = n + 1 + d$-dimensional space-time was suggested. The generalized Newton's law force reads

$$F = G\frac{m_1 m_2}{R^d} cos^2\theta, \tag{1.1}$$

where G is N-dimensional gravitational constant, m_1 and m_2 are masses of interacting point-like particles and R is the distance between them (that is supposed to be large enough). Here θ is the angle between the direction of motion of these two particles in n-dimensional time manifold $\mathbb{R}^n$, or, more explicitly,

$$\cos\theta = \vec{n}_1 \vec{n}_2, \tag{1.2}$$

where n-dimensional unit vectors $\vec{n}_1$, $\vec{n}_1$ define the motion of two particles in the time manifold $\mathbb{R}^n$

$$\vec{t}_1(\tau) = \vec{n}_1 \tau + \vec{t}_{01}, \quad \vec{t}_2(\tau) = \vec{n}_2 \tau + \vec{t}_{02}. \tag{1.3}$$

---

[1] ivashchuk@mail.ru

2 The "derivation"

The formula (1.1) may be obtained using the following relation for gravitational potential energy

$$V = -\bar{G}\frac{tr(M_1 M_2)}{R^{d-1}}, \tag{2.1}$$

where $\bar{G} = G/(d-1)$, $M_1 = (m_1 n_1^i n_1^j)$ and $M_2 = (m_2 n_2^i n_2^j)$ are (multitemporal) mass matrices (inertial or gravitational ones) for the first and the second particles, respectively. (For $d = 2$ case see [3].)

Relation (2.1) is a special case of multitemporal analogue of Newton's law obtained in [2]

$$V = -\bar{G}\frac{tr(M_{1G} M_{2I})}{R^{d-1}}, \tag{2.2}$$

where M_{1G} is a gravitational mass matrix of the first particle and M_{2I} an inertial mass matrix of the second particle.

The formula (2.2) was derived in [2] by considering the motion of the relativistic massive particle in a gravitational background described by multitemporal generalization of the Tangherlini solution [5, 6] (see [2] and Appendix). The derivation of (2.2) was performed in assumption of a non-relativistic motion of the particle at large distances from the gravitating center. Recall that the inertial mass matrix for the relativistic particle of scalar mass m is $M = (mn^i n^j)$, where the unit vector $\vec{n} = (n^i)$ describes the motion in the time manifold $\mathbb{R}^n$: $\vec{t}(\tau) = \vec{n}\tau + \vec{t}_0$, where τ is a synchronous time parameter.

The solution in [2] contains both black hole configurations and naked singularities as well. The relation for M_{1G} in [2] contains some extra parameters, but for black hole configurations in a space-time with extra time dimensions (that are trivial generalization of Tangherlini solution [5]) we get that the gravitational and inertial mass matrices are coinciding, i.e.

$$M_{iG} = M_{iI}, \tag{2.3}$$

$i = 1, 2$ (see Appendix).

3 Discussions

The relation (1.1) implies the following value for the effective gravitational constant

$$G_{eff} = Gcos^2\theta, \tag{3.1}$$

where $\theta = \theta_{ij}$ refers to a pair of two particles labelled by indices i and j.

It follows from (3.1) that $G_{eff} = 0$ for $\theta = \pi/2$, i.e. the "Newtonian" interaction is absent in this case. It is important that

$$G_{eff} \geq 0, \tag{3.2}$$

i.e. the repulsion is absent.

Now we put $d = 2$. Let us consider a flow of particles moving in the time manifold $\mathbb{R}^n$ near the t^1-axis with small enough values of $\theta_{ij} = \theta_{ij}(t^1)$. Then, for proper restrictions on θ_{ij} and $\dot{\theta}_{ij}$ imposed, the effective 4-dimensional gravitational constant G_{eff} may have a small enough variation in t^1-time (compatible with the observational data) and the weak equivalence principle will be not violated.

Acknowledgments

This work was supported in part by the Russian Foundation for Basic Research grants Nr. 09-02-00677-a.

Appendix

The solution in [2] reads

$$\begin{aligned} g = \quad & -\sum_{i=1}^{n}(1 - BR^{1-d})^{a_i} dt^i \otimes dt^i \\ & +(1 - BR^{1-d})^{b-1} dR \otimes dR + (1 - BR^{1-d})^{b} R^2 d\Omega^2_d, \end{aligned} \tag{A.1}$$

where

$$b = (1 - \sum_{i=1}^{n} a_i)/(d - 1) \tag{A.2}$$

and the parameters $a_1, \ldots, a_n$ satisfy the relations

$$(\sum_{i=1}^{n} a_i)^2 + (d-1)\sum_{i=1}^{n} a_i^2 = d. \tag{A.3}$$

Here B is constant. This solution is a special case of the solution obtained earlier in [4].

The black hole configuration takes place when the set $(a_1, \dots, a_n)$ is coinciding with one of the points

$$(1, 0, \dots, 0), \dots, (0, \dots, 0, 1). \tag{A.4}$$

In other cases we get naked singularities [2].

The gravitational mass matrix corresponding to (A.1) was defined in [2] as follows

$$M_{ij} = a_i \frac{\delta_{ij} B}{2G}. \tag{A.5}$$

More general form of solution may be obtained by a $O(n)$-rotation of time axes $t^1, ..., t^n$. For the set $(1, 0, \dots, 0)$ we get a "pure state" (projector) matrix in (A.5) which may be reduced to the general form $M = (mn^i n^j)$ (where $\vec{n} = (n^i)$ is a vector of unit length) by a suitable $O(n)$-rotation.

References

[1] V. D. Ivashchuk, On multitemporal anlogue of Newton's gravitational law and variation of gravitational constant, A talk given at seminar of RGS-VNIIMS, November, 2007.

[2] V.D. Ivashchuk and V.N. Melnikov, Multi-temporal Generalization of the Tangherlini Solution, *Class. Quantum Grav.* **11**, (1994) 1793-1805.

[3] V.D. Ivashchuk and V.N. Melnikov, Multitemporal Generalization of the Schwarzschild Solution, *Int. J. Mod. Phys.*, **D 4**, No 2, (1995) 167-173.

[4] S.B. Fadeev, V.D. Ivashchuk and V.N. Melnikov, On Black Holes in Multidimensional Theory with Ricci-flat Internal Spaces, *Phys. Lett.*, **A 161** (1991) 98-100.

[5] F.R. Tangherlini, *Nuovo Cimento* **27** (1963) 636.

[6] R.C. Myers and M.J. Perry, *Ann. of Phys.* **172** (1986) 304.